

Understanding the Health Risks of E-Cigarettes



We have much more to learn about e-cigarettes. According to the Surgeon General, the CDC, and the National Academies of Science, Engineering and Medicine (NASEM), e-cigarettes deliver fewer toxins than combustible cigarettes, but that doesn't mean that they are safe or without risk.¹ In a 2020 report, the Surgeon General found that "the long-term health effects of using these products remain unknown, and short-term risks are only slowly coming into focus."²

E-Cigarettes are Not Harmless Water Vapor

E-cigarette aerosol can contain nicotine, cancer-causing chemicals and other toxins.

- While aerosol from e-cigarettes, including disposable e-cigarettes, contains fewer of most of the chemicals and toxins compared to cigarette smoke, it can still contain harmful and potentially harmful chemicals, including nicotine, formaldehyde, acrolein, volatile organic compounds like toluene, synthetic cooling agents, tobacco-specific nitrosamines, and metals like nickel and lead.³ Researchers at Johns Hopkins University identified a complex mixture of approximately 2,000 chemicals in four e-cigarette brands. The vast majority of these chemicals were unidentified.⁴ More research is needed about the chemicals found in e-cigarettes and the impact of inhaling these chemicals deeply into the lungs.⁵
- Flavored e-cigarettes may pose unique harms. CDC warns that the e-cigarette aerosol inhaled by users can contain harmful or potentially harmful chemicals, including "[f]lavorings such as diacetyl, a chemical linked to a serious lung disease. Some flavorings used in e-cigarettes may be safe to eat but not to inhale because the lungs process substances differently than the gut."⁶ In its 2024 report, the Surgeon General concluded, "Determinations that flavorants are 'generally recognized as safe' for use in foods are not applicable to the inhalation of tobacco products. These ingredients, when inhaled, may be directly toxic to the lungs or could result in higher absorption of toxicants."⁷ FDA has previously stated, "Flavorings that are safe for use in food may become toxic when these chemicals are heated and inhaled. Some have been shown to be harmful to the lungs."⁸

Emerging Evidence of the Harmful Effects of E-cigarettes

- Little is known about the long-term effects of e-cigarette use. Because the popularity of these products is relatively new and users have not had sufficient time to develop symptoms, there is not enough information to assess the long-term impact on cancer, respiratory disease, and heart disease risk.⁹
- In 2021, the WHO Global Tobacco Control Report detailed the emerging evidence of the harmful effects of e-cigarettes, including studies suggesting that e-cigarettes have negative effects on aspects of cardiovascular and respiratory health.¹⁰
- A 2020 Surgeon General's report also noted that, "Several studies demonstrate e-cigarette aerosol contains fine and ultrafine particles, such that use of the products could potentially increase cardiovascular and respiratory risks."¹¹
- E-cigarettes have been found to increase heart rate and blood pressure.¹² A 2021 report of the World Heart Federation identified other concerning changes related to e-cigarette use. The report concluded that more research on the long term effects of e-cigarettes is needed, but the "effects on the cardiovascular system could be substantial."¹³ A more recent scientific statement from the American Heart Association (AHA) on the cardiopulmonary impact of electronic cigarettes and vaping products concluded that "The presence of any acute cardiovascular effects of [e-cigarettes]

Understanding the Health Risks of E-Cigarettes

suggests that [e-cigarette] use is not benign and raises the possibility that adverse impact could accumulate over time.”¹⁴

- There is growing evidence that vaping can harm lung health. E-cigarette aerosol inhibits several kinds of immune cells in the lungs, compromising the ability to fight infection.¹⁵ Initial evidence also indicates that e-cigarette aerosols promote inflammation in the respiratory system.¹⁶ Additionally, nicotine, a prominent component of e-cigarette aerosol, is known to suppress immune function throughout the body.¹⁷
- An extensive review of the evidence on the effects of e-cigarettes on respiratory health noted that the development of serious lung disease usually follows decades of chronic smoking, “so the population effects of e-cigarette use may not be apparent until the middle of this century.” The researchers conclude that “current knowledge of these effects is insufficient to determine whether the respiratory health effects of e-cigarette are less than those of combustible tobacco products.” Based on the emerging evidence, they found that “e-cigarettes will likely prove to have at least some pulmonary toxicity with chronic and possibly even short-term use.”¹⁸
- Initial research indicates that e-cigarette aerosol can damage DNA.¹⁹ Recently, one study found similarly increased levels of DNA damage among exclusive e-cigarette users and exclusive cigarette smokers, compared to nonusers.²⁰ As noted by the Surgeon General’s Report and a recent policy statement from the American Association for Cancer Research (AACR) and the American Society of Clinical Oncology (ASCO), it is biologically plausible that over time the DNA damage from long-term exposure to e-cigarette aerosols could increase risk of cancer. However, the Surgeon General, AACR, and ASCO also agree that the evidence remains uncertain whether the level of exposure is high enough to contribute to the formation of cancer in humans.²¹
- CDC warns that e-cigarettes and other products containing nicotine are not safe to use during pregnancy, noting that “Nicotine is a health danger for a pregnant woman and her developing fetus. It can damage the brain and lungs as they develop.”²²
- Emerging research suggests that e-cigarette use has a negative effect on oral health.²³ Nicotine impacts blood flow to the gums and could lead to inflammation of the gums and gum disease after continued use.²⁴ The chemicals in e-cigarettes also stay in the mouth, and can soften tooth enamel and contribute to tooth decay.²⁵ Vaping also changes the balance of bacteria in the mouth in ways that put users at higher risk of oral infections.²⁶
- The variation in e-cigarettes now on the market can also impact their potential health impacts. Devices can have powerful batteries that create more aerosol and expose users to more toxicants. The proliferation of e-liquids with nicotine salts allow users to inhale significantly higher levels of nicotine.²⁷

Any Potential Benefit To Smokers Comes From Complete Switching, Yet Many E-Cigarette Users Continue To Smoke

- “Lower risk” or “less harmful” is only relevant when you are talking about an adult cigarette smoker switching completely to an e-cigarette. According to the 2022 NHIS, the majority of adults using e-cigarettes either still smoke (dual use 26.7%) or have never been smokers (32.2%).²⁸
- While noting that the evidence is insufficient to recommend e-cigarettes for smoking cessation, CDC states that, “E-cigarettes may have the potential to benefit adults who smoke and are not pregnant if they are used as a complete substitute for smoked tobacco products.”²⁹ Otherwise, dual use, even with cutting back the number of cigarettes smoked, still elevates smokers’ health risks for things like cardiovascular disease.³⁰ According to the NASEM report, “there is no available evidence whether or not long-term e-cigarette use among smokers (dual use) changes morbidity or mortality compared with those who only smoke combustible tobacco cigarettes.”³¹

E-Cigarettes Pose Significant Risks to Kids' Health

- According to the Surgeon General, “E-cigarette use poses a significant – and avoidable – health risk to young people in the United States. Besides increasing the possibility of addiction and long-term harm to brain development and respiratory health, e-cigarette use is associated with the use of other tobacco products that can do even more damage to the body.”³²
- The Surgeon General concluded that, “The use of products containing nicotine in any form among youth, including in e-cigarettes, is unsafe.”³³
- Similarly, the FDA warns that “Vaping is not harmless. ... Teens who vape may experience coughing, wheezing, nausea, vomiting, headaches, and dizziness.”³⁴
- While it’s too early to confirm long-term health outcomes – “it took ~50 years before the connection between tobacco smoking and the multitude of smoking-related diseases became clear” - AHA’s 2022 scientific statement, focused on the cardiopulmonary consequences of vaping in adolescents, found that:³⁵
 - Available studies on short-term effects, such as changes in vascular stiffness, blood pressure, and heart rate, suggest that vaping during adolescence may lead to increased likelihood of cardiovascular disease later in life.
 - Further, “[l]ung development continues into the early 20s, and adolescents who vape are at risk for stunting or altering their lung development such that they never reach their full lung function potential, ... humans who vape during their adolescent years will likely have lower lung function and thus will become symptomatic at a lower threshold with any additional insult such as asthma, chronic obstructive pulmonary disease, pneumonia, or interstitial lung disease.”
- In 2021, feelings of anxiety, stress, and depression was reported as the most common reason for continuing to use e-cigarettes among current youth e-cigarette users.³⁶ Early research has found that youth e-cigarette use is associated with mental health symptoms, such as depression.³⁷ When youth depend on nicotine, the addiction can lead to a cycle of temporary symptoms of nicotine withdrawal, such as irritability, restlessness, and feelings of anxiety and depression.³⁸ This cycle can be yet another source of stress.
- Nicotine is a highly addictive drug that can have lasting damaging effects on adolescent brain development and impacts attention, memory and learning.³⁹ According to the Surgeon General, nicotine can also prime the brain for addiction to other drugs.⁴⁰
- Some of the pod systems and disposable products deliver massive doses of nicotine, putting youth users at greater risk of addiction. Juul has stated that each Juul “pod” (cartridge of nicotine) delivers as much nicotine as a pack of 20 cigarettes. Many e-cigarette products use nicotine salts, which allow users to inhale high levels of nicotine more easily and with less irritation. As a result, it could be easier for young people to initiate the use of and quickly become addicted to nicotine with these products.⁴¹
- Studies have found that young people who use e-cigarettes are more likely to become smokers, and many are low-risk youth who would not have otherwise smoked cigarettes.⁴² Both the U.S. Surgeon General and NASEM have concluded that there is an association between e-cigarette use and subsequent cigarette smoking initiation.
 - In 2016, the Surgeon General concluded that e-cigarette use is “strongly associated” with the use of other tobacco products among youth and young adults, including conventional cigarettes.⁴³

Understanding the Health Risks of E-Cigarettes

- A 2018 report by the National Academies of Sciences, Engineering and Medicine (NASEM) concluded that “There is substantial evidence that e-cigarette use increases risk of ever using combustible tobacco cigarettes among youth and young adults.”⁴⁴
- One study found that youth who used e-cigarettes were four times more likely to subsequently try cigarettes.⁴⁵
- The concept of “lower risk” absolutely does not apply to youth – any nicotine use by youth is a serious concern.

The Claim that E-Cigarettes are 95% Safer than Cigarettes: Erroneous and Widely Disputed by Researchers

- This specific claim has serious limitations and has received criticism from different scientific sources:
 - A 2020 AJPH article conclusively refuted the erroneous claim that e-cigarettes have been proven to be 95% safer than regular cigarettes, concluding, “The “95% safer” estimate is a “factoid”: unreliable information repeated so often that it becomes accepted as fact.”
 - This article notes “the evidence-lacking estimate derived in 2013 cannot be valid today and should not be relied upon further.” Since 2013, a substantial amount of new evidence has emerged about e-cigarettes. The article notes that the devices are now more powerful, create more aerosol, and expose users to more toxicants. The proliferation of e-liquids with nicotine salts allow users to inhale significantly higher levels of nicotine. More research emerged about the toxicants in e-cigarettes, and their potential respiratory and cardiovascular effects.⁴⁶
 - As an editorial from The Lancet summarized, “the opinions of a small group of individuals with no pre-specified expertise in tobacco control were based on an almost total absence of evidence of harm. It is on this extraordinarily flimsy foundation that PHE based the major conclusion and message of its report.”⁴⁷
- The authors of the original estimate acknowledge they were limited by a “lack of hard evidence for the harms of most products on most of the criteria.” The introduction to the article noted that, “our understanding of the potential hazards associated with using electronic nicotine delivery systems (ENDS, e.g. E-cigarettes) is at a very early stage.”⁴⁸
- Given that studies continue to raise new concerns about e-cigarettes, the evidence is insufficient to reach definitive conclusions regarding their relative health risk compared to cigarettes. Recent evidence suggests that some risk reductions associated with e-cigarettes may not be as substantial as previously claimed.⁴⁹

Conclusion

While the evidence to date indicates that e-cigarettes expose users to fewer harmful chemicals than traditional cigarettes, the evidence of the harms of these products continues to emerge and the growing body of evidence indicates that they are certainly not safe, especially for kids and others who do not currently use tobacco products.

¹ National Academies of Sciences, Engineering, and Medicine (NASEM), Public Health Consequences of E-Cigarettes, Washington, DC: The National Academies Press, 2018, <http://nationalacademies.org/hmd/Reports/2018/public-health-consequences-of-e-cigarettes.aspx>. CDC, Health Effects of Vaping, May 15, 2024, accessed November 22, 2024 at <https://www.cdc.gov/tobacco/e-cigarettes/health-effects.html>.

² U.S. Department of Health and Human Services (HHS). Smoking Cessation. A Report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2020.

- ³ Kopa-Stojak PN, Pawliczak R. Disposable electronic cigarettes—chemical composition and health effects of their use. A systematic review. *Toxicology Mechanisms and Methods*. 2024 Nov 8;1-2.; Cheng, T, “Chemical Evaluation of Electronic Cigarettes,” *Tobacco Control* 23:iii1-iii17, May 2014, http://tobaccocontrol.bmj.com/content/23/suppl_2/iii1.full. Goniewicz, ML, et al., “Levels of selected carcinogens and toxicants in vapour from electronic cigarettes,” *Tobacco Control* 23(2):133-9, March 6, 2013. Williams, M, et al., “Metal and Silicate Particles Including Nanoparticles Are Present in Electronic Cigarette Cartomizer Fluid and Aerosol,” *PlosOne*, 8(3), March 2013. See also Williams, M, “Electronic Cigarette Liquids and Vapors: Is It Harmless Water Vapor,” presented October 3, 2013 at TRDRP Electronic Cigarette Webinar, <http://www.trdrp.org/docs/Williams%20ecig%20vapor%20this%20time%20slides%202013.pdf>. NASEM, Public Health Consequences of E-Cigarettes, 2018.
- ⁴ Tehrani MW, Newmeyer MN, Rule AM, Prasse C. Characterizing the Chemical Landscape in Commercial E-Cigarette Liquids and Aerosols by Liquid Chromatography-High-Resolution Mass Spectrometry. *Chemical Research in Toxicology*. 2021 Oct 5.
- ⁵ NASEM, Public Health Consequences of E-Cigarettes, 2018.
- ⁶ CDC, Health Effects of E-Cigarettes, May 15, 2024, accessed November 22, 2024 at <https://www.cdc.gov/tobacco/e-cigarettes/health-effects.html>.
- ⁷ HHS, Eliminating Tobacco-Related Disease and Death: Addressing Disparities: A Report of the Surgeon General. Atlanta, GA: HHS, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2024.
- ⁸ Gottlieb, S and Abernethy, A. Understanding the Health Impact and Dangers of Smoke and ‘Vapor’. FDA Voices: Perspectives from FDA Leadership and Experts. <https://web.archive.org/web/20190914195650/https://www.fda.gov/news-events/fda-voices-perspectives-fda-leadership-and-experts/understanding-health-impact-and-dangers-smoke-and-vapor>.
- ⁹ Rose JJ, et al. Cardiopulmonary Impact of Electronic Cigarettes and Vaping Products: A Scientific Statement From the American Heart Association. *Circulation*. 2023 Jul 17.; Siddiqi TJ, et al. Association of Electronic Cigarette Exposure on Cardiovascular Health: A Systematic Review and Meta-Analysis. *Current Problems in Cardiology*. 2023 Apr 22:101748.; Herbst RS, Hatsukami D, et al. Electronic Nicotine Delivery Systems: An Updated Policy Statement from the American Association for Cancer Research and the American Society of Clinical Oncology. *Clinical Cancer Research*. 2022 Nov 14;28(22):4861-70.; Kundu, A., et al. (2025). Evidence update on the cancer risk of vaping e-cigarettes: A systematic review. *Tobacco Induced Diseases*, 23(January), 6. <https://doi.org/10.18332/tid/192934>; NASEM, Public Health Consequences of E-Cigarettes, 2018.
- ¹⁰ WHO report on the global tobacco epidemic 2021: addressing new and emerging products. Geneva: World Health Organization; 2021. <https://www.who.int/publications/i/item/9789240032095>
- ¹¹ HHS. Smoking Cessation. A Report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2020.
- ¹² NASEM, Public Health Consequences of E-Cigarettes, 2018.
- ¹³ Bianco E, et al. E-Cigarettes: A New Threat to Cardiovascular Health – A World Heart Federation Policy Brief. *Global Heart*. Oct 2021; 16(1): 72. DOI: <https://doi.org/10.5334/gh.1076>
- ¹⁴ Rose JJ, et al. Cardiopulmonary Impact of Electronic Cigarettes and Vaping Products: A Scientific Statement From the American Heart Association. *Circulation*. 2023 Jul 17.
- ¹⁵ Gotts, JE, et al., “What are the respiratory effects of e-cigarettes?” *British Medical Journal* 366:l5275, 2019, <https://www.bmj.com/content/bmj/366/bmj.l5275.full.pdf>.
- ¹⁶ Traboulsi, Hussein et al. “Inhalation Toxicology of Vaping Products and Implications for Pulmonary Health.” *International journal of molecular sciences* vol. 21,10 3495. 15 May 2020, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7278963/>.
- ¹⁷ 2014 SGR, at 117.
- ¹⁸ Gotts, J. et al., “What are the respiratory effects of e-cigarettes?” *The BMJ*, 366:l5275, published online September 30, 2019. <https://www.bmj.com/content/bmj/366/bmj.l5275.full.pdf>
- ¹⁹ NASEM, Public Health Consequences of E-Cigarettes, 2018.
- ²⁰ Tommasi S, et al. Vaping Dose, Device Type, and E-Liquid Flavor are Determinants of DNA Damage in Electronic Cigarette Users. *Nicotine and Tobacco Research*. 2023 Feb 14:ntad003.
- ²¹ HHS. Smoking Cessation. A Report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2020.; Herbst RS, Hatsukami D, et al. Electronic Nicotine Delivery Systems: An Updated Policy Statement from the American Association for Cancer Research and the American Society of Clinical Oncology. *Clinical Cancer Research*. 2022 Nov 14;28(22):4861-70.
- ²² CDC, “E-Cigarettes and Pregnancy,” May 15, 2024, accessed December 24, 2024 at <https://www.cdc.gov/maternal-infant-health/pregnancy-substance-abuse/e-cigarettes.html>.
- ²³ Charde P, Ali K, Hamdan N. Effects of e-cigarette smoking on periodontal health: A scoping review. *PLOS Global Public Health*. 2024 Mar 20;4(3):e0002311.
- ²⁴ Kumar, P., Geisinger, M., DeLong, H. R., Lipman, R. D., & Araujo, M. W. (2020). Living under a cloud: Electronic cigarettes and the dental patient. *The Journal of the American Dental Association*, 151(3), 155-158. <https://jada.ada.org/action/showPdf?pii=S0002-8177%2820%2930004-0>.
- ²⁵ Yang, I., et al. (2020). The oral health impact of electronic cigarette use: a systematic review. *Critical Reviews in Toxicology*, 50(2), 97-127. <https://www.tandfonline.com/doi/abs/10.1080/10408444.2020.1713726>.
- ²⁶ Pushalkar, S., et al. (2020). Electronic Cigarette Aerosol Modulates the Oral Microbiome and Increases Risk of Infection. *Iscience*, 100884.; Ganesan, S. M., et al. (2020). Adverse effects of electronic cigarettes on the disease-naïve oral microbiome. *Science Advances*, 6(22), eaaz0108. <https://advances.sciencemag.org/content/6/22/eaaz0108>
- ²⁷ Eissenberg, T., et al., “Invalidity of an Oft-Cited Estimate of the Relative Harms of Electronic Cigarettes,” *AJPH*, published online January 8, 2020, <https://aiph.aphapublications.org/doi/10.2105/AJPH.2019.305424>.
- ²⁸ CDC, “Tobacco Product Use Among Adults—United States, 2022,” September 10, 2024, <https://www.cdc.gov/tobacco/media/pdfs/2024/09/cdc-osh-ncis-data-report-508.pdf>.
- ²⁹ CDC, “Vaping and Quitting,” May 15, 2024, accessed December 24, 2024 at <https://www.cdc.gov/tobacco/e-cigarettes/quitting.html>.
- ³⁰ Tverdal, A and Bjartveit, K, “Health Consequences of Smoking 1-4 Cigarettes per Day,” *Tobacco control* 14(5), 2005. Hackshaw, A, et al., “Low cigarette consumption and risk of coronary heart disease and stroke: meta-analysis of 141 cohort studies in 55 study reports,” *BMJ* 360:j5855, <http://doi.org/10.1136/bmj.j5855>, 2018. Hackshaw, A, et al., “Low cigarette consumption and risk of coronary heart disease and stroke: meta-analysis of 141 cohort studies in 55 study reports,” *BMJ* 360:j5855, <http://doi.org/10.1136/bmj.j5855>, 2018. HHS, The Health Consequences of Smoking: A Report of the Surgeon General, CDC, OSH, 2004. HHS, How Tobacco Smoke Causes Disease: The Biology and Behavioral Basis for Smoking-Attributable Disease, CDC, OSH, 2010. HHS, Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General, CDC, OSH, 2012, <https://www.ncbi.nlm.nih.gov/books/NBK99237/>. HHS, The Health Consequences of Smoking: 50 Years of Progress. A Report of the Surgeon General, CDC, OSH, 2014. Schane, RE, Ling, PM, & Glantz, SA, “Health Effects of

- Light and Intermittent Smoking: A Review,” *Circulation* 121(3):1518-1522, 2010. Tverdal, A & Bjartveit, K, “Health consequences of reduced daily cigarette consumption,” *Tobacco Control* 15:472-480, 2006.
- ³¹ NASEM, Public Health Consequences of E-Cigarettes, 2018.
- ³² HHS, Know the Risks: E-Cigarettes & Young People, web archive from May 10, 2024 accessed December 24, 2024 at <https://web.archive.org/web/20240510010628/https://e-cigarettes.surgeongeneral.gov/knowtherisks.html>.
- ³³ HHS, E-Cigarette Use Among Youth and Young Adults. A Report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2016.
- ³⁴ FDA CTP. “Resources for Professionals about Vaping & E-Cigarettes: a Toolkit for Working With Youth.” <https://digitalmedia.hhs.gov/tobacco/hosted/Vaping-E-Cigarettes-Youth-Toolkit.pdf>; See also Mukerjee, Richa, et al. “ENDS, cigarettes, and respiratory illness: Longitudinal associations among US youth.” *American Journal of Preventive Medicine* (2023).; Chaiton M, Pienkowski M, Musani I, Bondy SJ, Cohen JE, Dubray J, Eissenberg T, Kaufman P, Stanbrook M, Schwartz R. Smoking, e-cigarettes and the effect on respiratory symptoms among a population sample of youth: Retrospective cohort study. *Tobacco Induced Diseases*. 2023;21.
- ³⁵ Wold LE, et al, American Heart Association Council on Basic Cardiovascular Sciences; Council on Arteriosclerosis, Thrombosis and Vascular Biology; Council on Hypertension; and Stroke Council. Cardiopulmonary Consequences of Vaping in Adolescents: A Scientific Statement From the American Heart Association. *Circulation Research*. 2022 Jul 22;131(3):e70-82.
- ³⁶ CDC, Tobacco Product Use and Associated Factors Among Middle and High School Students — National Youth Tobacco Survey, United States, 2021. *MMWR Surveill Summ* 2022;71(No. SS-5):1-29.
- ³⁷ CDC, Health Effects of Vaping, May 15, 2024, accessed Dec 24, 2024 at <https://www.cdc.gov/tobacco/e-cigarettes/health-effects.html>.; Baiden P, et al. Use of electronic vaping products and mental health among adolescent high school students in the United States: The moderating effect of sex. *Journal of psychiatric research*. 2022 Mar 1;147:24-33.; Becker TD, Arnold MK, Ro V, Martin L, Rice TR. Systematic review of electronic cigarette use (vaping) and mental health comorbidity among adolescents and young adults. *Nicotine and Tobacco Research*. 2021 Mar;23(3):415-25.; Lechner WV, et al. Bi-directional associations of electronic and combustible cigarette use onset patterns with depressive symptoms in adolescents. *Preventive medicine*. 2017 Mar 1;96:73-8.
- ³⁸ U.S. Department of Health and Human Services. Smoking Cessation: A Report of the Surgeon General. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2020 [accessed 2022 April 4].
- ³⁹ HHS, The Health Consequences of Smoking: 50 Years of Progress. A Report of the Surgeon General, CDC, Office of Smoking and Health (OSH), 2014, https://www.ncbi.nlm.nih.gov/books/NBK179276/pdf/Bookshelf_NBK179276.pdf. See also: CDC, “Health Effects of Vaping,” May 15, 2024, accessed December 24, 2024, <https://www.cdc.gov/tobacco/e-cigarettes/health-effects.html>.
- ⁴⁰ HHS, E-Cigarette Use Among Youth and Young Adults. A Report of the Surgeon General—Fact Sheet. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2016.
- ⁴¹ Office of the Surgeon General, “Surgeon General’s Advisory on E-Cigarette Use Among Youth,” December 18, 2018, <https://stacks.cdc.gov/view/cdc/153187>.
- ⁴² Barrington-Trimis, JL, et al., “E-Cigarettes and Future Cigarette Use,” *Pediatrics*, 138(1), published online July 2016. Wills, TA, et al., “E-cigarette use is differentially related to smoking onset among lower risk adolescents,” *Tobacco Control*, published online August 19, 2016. Berry, KM, et al., “Association of Electronic Cigarette Use with Subsequent Initiation of Tobacco Cigarettes in US Youths,” *JAMA Network Open*, 2(2), published online February 1, 2019.
- ⁴³ HHS, E-Cigarette Use Among Youth and Young Adults. A Report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2016.
- ⁴⁴ NASEM, Public Health Consequences of E-Cigarettes, 2018.
- ⁴⁵ Berry, KM, et al., “Association of Electronic Cigarette Use with Subsequent Initiation of Tobacco Cigarettes in US Youths,” *JAMA Network Open*, 2(2), published online February 1, 2019.
- ⁴⁶ Eissenberg, T., et al., “Invalidity of an Oft-Cited Estimate of the Relative Harms of Electronic Cigarettes,” *AJPH*, published online January 8, 2020, <https://ajph.aphapublications.org/doi/10.2105/AJPH.2019.305424>.
- ⁴⁷ “E-cigarettes: Public Health England’s evidence-based confusion,” *The Lancet*, August 29, 2015.
- ⁴⁸ Nutt, DJ, et al., “Estimating the Harms of Nicotine-Containing Products Using the MCDA Approach,” *European Addiction Research* 20:218-225, 2014.
- ⁴⁹ Glantz SA, et al. Population-based disease odds for e-cigarettes and dual use versus cigarettes. *NEJM evidence*. 2024 Feb 27;3(3):EVID0a2300229.